



Learn more at our website: VerdierEyeCenter.com

WHAT CAUSES KERATOCONUS?

Doctors do not know for sure why people have keratoconus. In some cases, it appears to be genetic (passed down in families). About 1 out of 10 people with keratoconus have a parent who has it too. Keratoconus is also associated with:

- Eye allergies
- Excessive eye rubbing, and
- Connective tissue disorders like Marfan syndrome and Ehlers-Danlos syndrome

WHAT ARE THE SYMPTOMS?

Keratoconus often starts when people are in their late teens to early 20s. The vision symptoms slowly get worse over a period of about 10 to 20 years.

Keratoconus often affects both eyes, and can lead to very different vision between the two eyes. Symptoms can differ in each eye, and they can change over time.

In the early stage, keratoconus symptoms can include:

- mild blurring of vision
- slightly distorted vision, where straight lines look bent or wavy
- increased sensitivity to light and glare
- increased nearsightedness or astigmatism (when your eye cannot focus as well as it should). As a result, you may need new eyeglass prescriptions often.


In later stages, keratoconus symptoms often include:

- more blurry and distorted vision
- not being able to wear contact lenses. They may no longer fit properly and they are uncomfortable.

Keratoconus usually takes years to go from early- to late-stage. For some people, though, keratoconus can get worse quickly. The cornea can swell suddenly and start to scar. When the cornea has scar tissue, it loses its smoothness and becomes less clear. As a result, vision grows even more distorted and blurry.

HOW IS KERATOCONUS DIAGNOSED?

A routine eye exam may cause a suspicion of keratoconus, but it is best diagnosed by special scans of your cornea. Your optometrist or ophthalmologist will review these scans for information about thickness and curvature to diagnose keratoconus and monitor for progression.



WHAT ARE THE TREATMENTS?

Keratoconus treatment depends on your symptoms. When your symptoms are mild, your vision can be corrected with eyeglasses. Later you may need to wear special hard contact lenses to help keep vision in proper focus.

Here are other ways that your ophthalmologist might treat keratoconus:

- **Corneal cross-linking (CXL).** Your ophthalmologist uses a special UV light and eye drops to strengthen the cornea. Doing this helps to flatten or stiffen your cornea, keeping it from bulging further.
- **Corneal transplant.** When symptoms are severe, your ophthalmologist may suggest a corneal transplant. Your ophthalmologist replaces all or part of your diseased cornea with healthy donor cornea tissue.
- **Scleral Contact Lens.** These are large-diameter lenses that provide clear and comfortable vision for keratoconus patients because the lenses do not actually touch the cornea.
- **Gas Permeable Contact Lens.** Rigid Gas Permeable lenses (also known as RGPs or simply GPs) are one of the most popular contact lenses for Keratoconus. These stable contact lenses essentially act as a second surface for the eye, compensating for the irregularities of the cornea and functioning as the eye's new reflective surface.

Do not rub your eyes!

With keratoconus, try to avoid rubbing your eyes. This can damage thin corneal tissue and make your symptoms worse.

If you have itchy eyes that cause you to rub, speak to your ophthalmologist about medicines to control your allergies.